



(11) Publication number:

0 192 492
A2

(13)

EUROPEAN PATENT APPLICATION

(21) Application number: 86301278.7

 (51) Int. Cl.⁴: **C 07 K 7/06**
A 61 K 37/02

(22) Date of filing: 21.02.86

(30) Priority: 22.02.85 US 704299

 (43) Date of publication of application:
 27.08.86 Bulletin 86/35

 (64) Designated Contracting States:
 AT BE CH DE FR GB IT LI LU NL SE

 (71) Applicant: **THE SALK INSTITUTE FOR BIOLOGICAL STUDIES**
 10010 North Torrey Pines Road
 La Jolla California 92037(US)

 (72) Inventor: **Rivier, Jean Edouard Frederic**
 9674 Blackgold Road
 La Jolla California 92037(US)

 (72) Inventor: **Anderson, Harry Alec**
 3525 Tony Drive
 San Jolla California 92122(US)

 (72) Inventor: **Wylie, Walker Vale, Jr.**
 1643 Valdez
 La Jolla California 92037(US)

 (74) Representative: **Lawrence, Malcolm Graham et al,**
Malcolm Lawrence & Co. 9th Floor Terminus House
Terminus Street
Harlow Essex CM20 1XF(GB)

(54) Peptides containing an aliphatic-aromatic ketone side chain.

(57) GnRH peptide analogs which regulate the secretion of gonadotropins by the pituitary gland and either promote or inhibit the release of steroids by the gonads. Administration of an effective amount of a GnRH antagonist prevents ovulation of female mammalian eggs and/or the release of gonadotropins. Administration of GnRH agonists can be used to regulate fertility in male and female mammals.

These and other peptide hormones exhibit improved binding efficiency and biological potency as a result having a residue in a critical, generally central location in the chain which residue contains a mixed alkyl ketone side-chain terminating in an aromatic group. Methods for efficiently synthesizing these peptides from readily available compounds are disclosed.

EP 0 192 492 A2

PEPTIDES CONTAINING AN ALIPHATIC-AROMATIC
KETONE SIDE CHAIN

0192492

The present invention relates to peptides which affect the release of gonadotropins or inhibit the release of GH by the pituitary gland in mammals, and to methods of making such peptides. More particularly, the present invention is directed to peptides which have improved biological potency to either promote or inhibit gonadal function and the release of the steroidal hormones, progesterone and testosterone or improved biological potency to inhibit the release of GH.

The pituitary gland is attached by a stalk to the region in the base of the brain known as the hypothalamus. In particular, follicle stimulating hormone (FSH) and luteinizing hormone (LH), sometimes referred to as gonadotropins or gonadotropic hormones, are released by the pituitary gland. These hormones, in combination, regulate the functioning of the gonads to produce testosterone in the testes and progesterone and estrogen in the ovaries, and they also regulate the production and maturation of gametes. Growth hormone (GH) is also released by the pituitary gland.

The release of a hormone by the anterior lobe of the pituitary gland usually requires a prior release of another class of hormones produced by the hypothalamus. One of the hypothalamic hormones acts as a factor that triggers the release of the gonadotropic hormones, particularly LH, and this hormone is referred to herein as GnRH although it has also been referred to as LH-RH and as LRF. GnRH has been isolated and characterized as a decapeptide having the following structure: pGlu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH₂.

Peptides are compounds which contain two or more amino acids in which the carboxyl group of one acid is linked to the amino group of the other acid. The

formula for GnRH, as represented above, is in accordance with conventional representation of peptides where the amino or N-terminus appears to the left and the carboxyl or C-terminus to the right. The position of the amino acid residue is identified by numbering the amino acid residues from left to right. In the case of GnRH, the hydroxyl portion of the carboxyl group of glycine has been replaced with an amino group(NH₂). The abbreviations for the individual amino acid residues above are conventional and are based on the trivial name of the amino acid, e.g. pGlu is pyroglutamic acid, His is histidine, Trp is tryptophan, Ser is serine, Tyr is tyrosine, Gly is glycine, Leu is leucine, Orn is ornithine, Arg is arginine, Lys is lysine, Cys is cysteine, Asn is asparagine, Thr is threonine, Pro is proline, Phe is phenylalanine, Glu is glutamic acid, Asp is aspartic acid and Ala is alanine. Except for glycine, amino acids of the peptides of the invention are of the L-configuration unless noted otherwise.

The substitution of a D-amino acid for Gly in the 6-position of the GnRH decapeptide or nonapeptide provides a GnRH analog having substantially greater binding affinity and thus can be used to produce both agonists and antagonists of higher potency. The substitution of an ethylamide moiety or the like for Gly-NH₂ at the C-terminus produces agonists of higher potency. Other substitutions throughout the GnRH decapeptide are known which produce antagonists having an inhibitory effect on the release of LH and other gonadotropins by the pituitary gland of mammals. Such a releasing or inhibitory effect is obtained when the GnRH analog is administered to a mammalian intravenously, subcutaneously, intramuscularly, orally, percutaneously, e.g. intranasally, intravaginally, or in delayed or timed-release formulations.

There are reasons for desiring to prevent ovulation in female mammals, and the administration

of GnRH analogs that are antagonistic to the normal function of GnRH or of large doses of agonists of GnRH have been used to suppress or delay ovulation. For this reason, such analogs of GnRH are being investigated for
5 their potential use as a contraceptive or for regulating conception periods. GnRH antagonists may also be used for the treatment of precocious puberty and endometriosis. Such antagonists have also been found useful to regulate the secretion of gonadotropins in
10 male mammals and can be employed to arrest spermatogenesis, e.g. as male contraceptives, and for treatment of prostatic hypertrophy.

It is desired to provide improved peptides which are more potent analogs of GnRH.

15

The present invention provides improved GnRH analogs some of which are strongly antagonistic to GnRH and have an inhibitory effect on the reproduction processes of mammals and others which are potent
20 agonists of GnRH.

Generally, in accordance with the present invention, GnRH peptides have been synthesized which have stronger binding efficiency as a result of including a residue having an aliphatic-aromatic ketone
25 side chain in the 6-position, but which can be synthesized economically using relatively inexpensive materials. The GnRH antagonists strongly inhibit the secretion of gonadotropins by the pituitary gland of mammals, including humans, and/or inhibit the release
30 of steroids by the gonads. These peptides are analogs of GnRH wherein a D-isomer alpha-amino acid having a carboxyl-containing side chain, e.g. D-Glu, D-Hgl or D-Asp, is originally located in the 6-position; the side-chain carboxyl group of this residue is then
35 converted to a mixed alkyl ketone via the formation of a side chain acylium ion intermediate from the carboxyl group, which occurs upon treatment with HF or an

equivalent acid, such as a suitable Lewis acid. This mixed alkyl ketone side chain should be one which terminates with an aromatic moiety. By mixed alkyl ketone, for purposes of this application, is meant a
5 ketone which contains one alkyl group and one non-alkyl group. By aromatic, for purposes of this application, is meant a resonant carbocyclic or heterocyclic group, such as that derived from anisole, indole, furan, alkyl pyrrole or thiophene. By Hgl is meant alpha-amino adipic
10 acid, which is also referred to as homoglutamic acid.

In addition, the GnRH antagonists include a 1-position substitution, such as D-pGlu, dehydro-Pro, Pro, halogenated D-Phe, D-Trp or β -(2-naphthyl)-D-alanine(hereinafter β -D-2NAL), a substituted
15 (preferably halogenated) D-Phe in the 2-position, a 3-position substitution, an optional substitution of a diamino acid having not more than 5 carbon atoms in the 4-position, an optional substitution in the 5-position in the form of a halogenated L-Phe, a halogenated L-Tyr
20 or L-Arg and optional substitutions in the 7-and 10 positions. The 1-position substituent, except for D-pGlu, is preferably modified so that its alpha amino group contains an acyl group, such as formyl(For), acetyl(Ac), acrylyl(Acr), vinylacetyl(Vac) or
25 benzoyl(Bz), with Ac and Acr being preferred. Modified D-Trp in the 3-position provides increased antagonistic activity as a result of the specific modifications present in the indole ring. Single substitutions for hydrogen are made in either the 5- or 6-position, and
30 the substitutions are selected from chloro, fluoro, bromo, methyl, amino, methoxy and nitro, with chloro, fluoro and nitro being preferred. The indole nitrogen may also be acylated, e.g. with formyl (N^{in} For- or lFor-) or with acetyl. Another 3-position substituent
35 is D-PAL which stands for D-alanine which is substituted by pyridyl on the β -carbon atom with the linkage being to the 2-, 3- or 4-position on the pyridine ring, with

D-3PAL being preferred. As mentioned above, the substitutions in the 4-, 7- and 10-positions are generally considered to be optional. If substituted, the 10-position is preferably D-Ala-NH₂.

5 Because these peptides are highly potent to inhibit release of LH, they are referred to as GnRH antagonists. The peptides inhibit ovulation of female mammals when administered at very low levels at proestrous and are also effective to cause resorption of
10 fertilized eggs if administered shortly after conception. These peptides are also effective for the contraceptive treatment of male mammals.

 The improved GnRH agonists have a similar residue in the 6-position and may have optional
15 substitutions in the 1-position, preferably formyl Pro, and in the 10-position, preferably -NHCH₂CH₃ (-NH₂Et).

 Other peptide hormones that likewise incorporate such a residue having an aliphatic-aromatic ketone side-chain at a critical generally central
20 location, which results in increased binding affinity to the receptor for that hormone, can be synthesized.

 More specifically, the peptides of the present invention from the standpoint of a GnRH antagonist are
25 represented by the following Formula I: X-R₁-(W)D-Phe-R₃-R₄-R₅-R₆(V)-R₇-Arg-Pro-R₁₀ wherein X is hydrogen or an acyl group having 7 or less carbon atoms; R₁ is dehydro-Pro, Pro, D-pGlu, D-Phe, D-Trp or B-D-NAL; W is F, Cl, Cl₂, Br, NO₂ or C Me/Cl; R₃ is
30 D-PAL, D-Trp, (NⁱⁿFor)D-Trp or D-Trp which is substituted in the 5- or 6-position with NO₂, NH₂, OCH₃, F, Cl, Br or CH₃; R₄ is Ser, Orn, AAL or aBu; R₅ is Tyr, Arg, (3F)Phe, (2F)Phe, (3I)Tyr, (3CH₃)Phe, (2CH₃)Phe, (3Cl)Phe or (2Cl)Phe; R₆ is
35 D-Glu, D-Hgl or D-Asp; R₇ is Leu, NML, Nle or Nva; R₁₀ is Gly-NH₂, D-Ala-NH₂ or NH-Y, with Y being lower alkyl, cycloalkyl, fluoro lower alkyl or NHCONH₂.

where Q is H or lower alkyl; and V is an aromatic moiety portion of a ketone formed from the carboxylic group side chain of R₆ and a compound selected from Class 2' consisting of ethylbenzene, propylbenzene,

- 5 isopropylbenzene, butylbenzene, s-butylbenzene, isobutylbenzene, t-butylbenzene, amylbenzene, 1-methylbutylbenzene, 1-ethylpropylbenzene, 3-methylbutylbenzene, 1,1-dimethylpropylbenzene, hexylbenzene, heptylbenzene, 2-ethylhexylbenzene,
- 10 octylbenzene, nonylbenzene, decylbenzene, dodecylbenzene, tetradecylbenzene, hexadecylbenzene, octadecylbenzene, cyclopropylbenzene, cyclopentylbenzene, cyclohexylbenzene, (4-acyloxycyclohexyl)-benzene,
- 15 3-methyl-5-phenyl-cyclohex-2-enone, o-xylene, m-xylene, p-xylene, m-ethyltoluene, p-ethyltoluene, o-propyltoluene, m-propyltoluene, m-cymene, p-propyltoluene, p-cymene, m-s-butyltoluene, p-s-butyltoluene, m-t-butyltoluene, p-dodecyltoluene,
- 20 p-diethylbenzene, m-t-butylethylbenzene, p-t-butylethylbenzene, m-diisopropylbenzene, p-diisopropylbenzene, p-dibutylbenzene, p-di-s-butylbenzene, p-di-t-butylbenzene, p-di-(1-methylbutyl)benzene, indane(indan or
- 25 hydrindene), 5-methylindane, 6-t-butyl indane, 2-benzyl indane, dialkyl indane, trialkyl indane, tetraalkyl indane, pentaalkyl indane, hexaalkyl indane, heptaalkyl indane, 1-carboethoxy indane, tetralin, 6-methyl tetralin, 6-ethyl tetralin, 6-butyl tetralin, 6-hexyl
- 30 tetralin, 6-cyclohexyl tetralin, dialkyl tetralin, tetraalkyl tetralin, pentaalkyl tetralin, 7-ethyl-1-carboethoxymethyl tetralin, 2-phenyl tetralin, hemimellitene, pseudocumene, mesitylene, prehnitene, isodurene, durene, pentamethylbenzene, ethyl-1,4-xylene,
- 35 4-ethyl-1,3-xylene, 2-propyl-1,4-xylene, 2-isopropyl-1,4-xylene, 4-propyl-1,2-xylene, 4-propyl-1,3-xylene, 2-isobutyl-1,4-xylene,

- 5-6-butyl-1,3-xylene, 2-t-amyl-4-isopropyltoluene,
trialkylbenzene, 4-benzyl-1,3-xylene,
3,4,5,11-tetrahydroacenaphthene, ethylmesitylene,
hydrindacene, hydrophenanthrene, pentaalkylbenzene,
5 hydropyrene, hydroanthracene, diphenylmethane,
diphenylpropane, bibenzyl, 3,4-diphenylhexane,
 α , ω -diphenylalkanes, triphenylmethane, paracyclophanes,
phenylethylenes, chalkones, formamidotoluene,
phenylacetic acid, alkyl phenylacetate,
10 phenylacetonitrile, desoxybenzoin,
1-phenyl-2-nitroethane, 1-phenyl-2-acetamidoethane,
alkyl 3-phenylpropionate, 3-phenylpropionitrile,
phenylbenzoylalkanes, phenylchloroalkanes,
phenylnitroalkanes, alkylphenylbutyrates,
15 haloalkylbenzenes, phenol, phenyl acetate, phenyl
propionate, phenyl benzoate, alkylphenols, halophenols,
catechol, resorcinol, alkyl resorcinol, pyrogallol,
phloroglucinol, trihydroxytoluene,
trihydroxyisoamylbenzene, anisole, phenetole,
20 alkylphenylethers, alkyltolylethers, ethylanisole,
p-t-butylanisole, m-heptylanisole, p-cyclohexylanisole,
anisylhexanes, dimethylanisole, 2-ethyl-4-methylanisole,
5-methoxytetralin, isopropylmethylanisole,
hydrophenanthrene, trialkylanisole, fluoroanisole,
25 chloroanisole, chlorophenetole, bromoanisole,
bromophenetole, iodoanisole, alkylhaloanisoles,
dialkylhaloanisoles, dihaloanisoles,
dialkoxyhaloanisoles, guaiacol, resorcinol monomethyl
ether, hydroquinone monomethyl ether,
30 alkylhydroxyanisoles, dihydric phenolic dimethyl ethers,
polyhydric phenolic methyl ethers, diphenyl ether,
alkyldiphenylethers, chlorodiphenyl ethers, alkoxy
diphenyl ethers, dialkyldiphenyl ethers,
nitrophenylethers, thioanisole, thiophenetole, alkyl
35 phenyl sulfides, o-tolylthioethers, alkylthioanisoles,
chlorothioanisole, diphenyl sulfide, nitrodiphenyl
sulfide, 2-thiocresol, 3-methoxythiophenol,

- 3-ethoxythiophenol, diphenyl disulfide, acetanilide (AA),
alkyl AA, dialkyl AA, acetamidoindane,
acetamidotetralin, trimethyl AA, chloro AA,
chloro-4-methyl AA, alkoxy AA, N,N-diacylanilines,
5 nitrobromobenzene, nitrophenol, nitroanisole,
nitrophenetole, hydroxy-3-nitrotoluene, nitroresorcinol,
nitroanisole, hydroxy-4-nitroanisole, benzoic acid,
m-toluic acid, salicylic acid, alkyl salicylates,
alkylalkoxy benzoates, alkyl hydroxybenzoates,
10 dimethylacetophenone, trimethylacetophenone,
trimethylpropiophenone, methoxyacetophenone,
dihydroxyacetophenone, dihydroxypropiophenone,
benzophenone, dimethylbenzophenone,
dihydroxybenzophenone, biphenyl (BP), alkyl BPs, dialkyl
15 BPs, 9-10-dihydrophenanthrene, chloro BP, bromo BP,
hydroxy BP, methoxy BP, methoxy-chloro BP, acetyl BP,
nitro BP, chloroacetyl BP, diphenylbenzene,
1,4-terphenyl, 1,3,5-triphenylbenzene, fluorene (F),
benzyl F, methoxy F, carbomethoxy F, benzoyl F,
20 naphthalene, alkyl naphthalenes, halonaphthalenes,
naphthol, alkyl naphthol ethers, 2-naphthyl methyl
sulfide, naphthosultone, 1,8-naphthosultam, naphthalene
carboxylic acids, anthracene, alkyl anthracenes, halo
anthracenes, alkyl halo anthracenes, alkylalkoxy
25 anthracenes, anthrophenone, 9,9'-bianthryl,
phenanthrene (P), alkyl P, halo P, alkoxy P, acetoxo P,
hydroxy P, 3-acetamido P, pyrene, 2-methylpyrene,
1-benzoylpyrene, chrysene, 2-ethylchrysene,
6-benzylchrysene, triphenylene, perylene, fluoranthene,
30 biphenylene, furan, alkyl furan, benzofuran (BF), methyl
BF, ethyl BF, propyl BF, benzyl BF, phenyl BF, anisyl
BF, dihydrobenzofuran, dibenzofuran (DBF), ethyl DBF,
propyl DBF, bromo DBF, methoxy DBF, nitro DBF, xanthene,
1-hydroxy-9-oxoxanthene, thiophene, alkylthiophenes,
35 dialkylthiophenes, trialkylthiophenes,
2-benzylthiophene, halothiophenes, dihalothiophenes,
trihalothiophenes, alkylhalothiophenes,

halophenylthiophene, methyl phenylthiophene, dithienyl,
 dimethyldithienyl, terthienyl, benzo[b]thiophene (BT),
 methyl BT, methoxy BT, dibenzothiophene, alkyl pyrrole,
 dialkyl pyrrole, trialkyl pyrrole, dialkyl-carbomethoxy
 5 pyrroles, indole, 2-methylindole, 3-methylindole,
 1,2-dimethylindole, 1,2,3-trimethylindole,
 2,3,4,6-tetramethylindole, 2-phenylindole,
 2,3-dimethyl-1-acetylindole, tetrahydrocarbazole (THC),
 9-acetyl THC, 9-benzoyl THC, 6-halo-9-acetyl THC,
 10 carbazole, acetyl carbazole, alkyl carbazole, haloalkyl
 carbazole, benzoyl carbazole, acetylinoline,
 1-acyl-2,3-dimethyl-indolines, hexahydrocarbazoles,
 phenyl pyrazole, phenylalkyl pyrazole,
 1-phenyl-3-pyrazolin-5-one (PP), alkyl PP, dialkyl PP,
 15 2-imidazolone (IA), 4-methyl IA,
 2-oxo-2,3-dihydrobenzimidazole,
 imidazo[1,5-a]pyridine (IP), methyl IP, 5,7-dimethyl
 quinoline, hydroxy quinoline, methoxy quinoline,
 2-methylhydroxy quinoline, acyl tetrahydroquinoline,
 20 acridan, 10-acetyl acridan, 2-hydroxy-4-methylthiazole,
 10-ethyl phenoxazine, 10-acetyl phenoxazine,
 phenothiazine (PT), 10-alkyl PT, 3,10-dimethyl PT,
 10-acyl PT, 1,2-benzisoxazole (BIO), 3-methyl BIO,
 7-methoxy BIO and 3-phenyl-7-methoxy BIO.

25 From the standpoint of a GnRH agonist, the
 peptides are represented by the Formula IA: R_1 -His-
 Trp-Ser-Tyr- R_6 (V)-Leu-Arg-Pro- R_{10} , wherein R_6 and
 V are defined as set forth above and R_1 is pGlu or
 For-Pro and R_{10} is Gly-NH₂, D-Ala-NH₂ or
 30 substituted amide.

By β -D-NAL is meant the D-isomer of alanine
 which is substituted by naphthyl on the β -carbon atom,
 which may also be designated 3-D-NAL. Preferably
 β -D-2NAL is employed which means that the β -carbon atom
 35 is attached to naphthalene at the 2-position on the ring
 structure; however, β -D-1NAL may also be used. Dap
 represents α , β -diaminopropionic acid, which is also

termed β -aminoalanine, and by NML is meant $N^{\alpha}CH_3$ -L-Leu. By AAL is also meant β -amino-Ala and by aBu is meant α, γ diamino butyric acid, either of which or Orn can be present in the 4-position. When Ser is not present in the 4-position, dehydro Pro is preferably present in the 1-position. By $C^{\alpha}Me/Cl$ -D-Phe is meant D-Phe having its α -carbon methylated and being substituted by Cl in the para-position.

The term $R_6(V)$ in Formulas I and IA is used to define the D-amino acid residue in the main peptide chain having its side chain carboxyl group modified to form a mixed alkyl ketone. Preferably, the residue in the main chain is D-Glu; however, it may instead be D-Hgl or D-Asp.

The peptides of the present invention can be synthesized by classical solution synthesis or by a solid phase technique using a chloromethylated resin, a benzhydrylamine (BHA) resin, a methylbenzhydrylamine resin (MBHA), an N-alkyl amino methyl resin (NAAM) or any other suitable resin known in the art. When using classical synthesis, it may be advantageous to independently generate the ketone side chain prior to linking the amino acid to the adjacent residues in the peptide chain. Solid phase synthesis is conducted in a manner to stepwise add the amino acids in the chain in the manner set forth in detail in the U.S. Patent No. 4,211,693. Side-chain protecting groups, as are well known in the art, are preferably added to Ser, Tyr and Arg when present, as well as to certain of the substituents, and may optionally be added to Trp (unless acylated), before these amino acids are coupled to the chain being built upon the resin. When solid phase synthesis is used, the D-Glu, D-Hgl or D-Asp residue in the 6-position is preferably protected with Bzl(benzyl ester), 2,6-dichlorobenzyl(DCB), dinitrophenyl(Dnp), 1-hydroxy-benzotriazole benzyl ester(OHbt), 8-hydroxy-quinoline ester(OHq), p-nitrobenzyloxy(ONBzl),

phenylazophenyl or tertiary butoxy; such a synthesis provides the fully protected intermediate peptidoresin.

The intermediates of the invention with respect to a GnRH antagonist may be represented by Formula II:

- 5 $X^1-R_1-(W)D-Phe-R_3(X^2)-R_4(X^3)-R_5(X^4 \text{ or } X^6)-R_6(X^5)-R_7-Arg(X^6)-Pro-X^7$ wherein: X^1 is an α -amino protecting group of the type known to be useful in the art in the stepwise synthesis of polypeptides and when X in the desired peptide composition is a
- 10 particular acyl group, that group may be used as the protecting group. Among the classes of α -amino protecting groups covered by X^1 are (1) acyl-type protecting groups, such as formyl(For), trifluoroacetyl, phthalyl, p-toluenesulfonyl(Tos), benzoyl(Bz),
- 15 benzenesulfonyl, o-nitrophenylsulfonyl(Nps), tritylsulfonyl, o-nitrophenoxycarbonyl, acrylyl(Acr), chloroacetyl, acetyl(Ac) and γ -chlorobutyryl; (2) aromatic urethan-type protecting groups, e.g., benzyloxycarbonyl (Z), fluorenylmethyloxycarbonyl(FMOC),
- 20 and substituted benzyloxycarbonyl, such as p-chlorobenzyloxycarbonyl (ClZ), p-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl and p-methoxybenzyloxycarbonyl; (3) aliphatic urethan protecting groups, such as tertbutyloxycarbonyl(Boc), diisopropylmethoxycarbonyl,
- 25 isopropylloxycarbonyl, ethoxycarbonyl and allyloxycarbonyl; (4) cycloalkyl urethan-type protecting groups, such as cyclopentylloxycarbonyl, adamantylloxycarbonyl and cyclohexylloxycarbonyl; (5) thiourethan-type protecting groups, such as
- 30 phenylthiocarbonyl; (6) alkyl-type protecting groups, such as allyl(Aly), triphenyl-methyl(trityl) and benzyl(Bzl); (7) trialkylsilane groups, such as trimethylsilane. The preferred α -amino protecting group is Boc when X is hydrogen.
- 35 X^2 is hydrogen or a protecting group for the indole nitrogen of Trp, such as formyl or benzyl. In many syntheses, there is no need to protect the indole

NH of Trp; however X^2 is formyl when R_3 is ($N^{in}For$)D-Trp. There is no need to protect D-3PAL.

X^3 is hydrogen or a protecting group for the alcoholic hydroxyl group of Ser, such as one selected
5 from the group consisting of acetyl, benzoyl, tetrahydropyranyl, tert-butyl, trityl, benzyl and 2,6-dichlorobenzyl, with benzyl being preferred. Alternatively, when a substitution is made for Ser, X^3 may be a protecting group for a side chain amino group,
10 such as Tos, Z or ClZ.

X^4 is hydrogen or a protecting group for the phenolic hydroxyl group of Tyr, if Tyr is present, selected from the group consisting of tetrahydropyranyl, tert-butyl, trityl, benzyl, 2, 4-bromobenzyloxycarbonyl
15 and 2,6-dichlorobenzyl. 2,6-dichlorobenzyl is preferred.

X^5 is a protecting group for the side chain carboxyl group of D-Glu, D-Hgl or D-Asp, selected from the group consisting of Bzl(benzyl ester), 2,6-dichlorobenzyl(DCB), dinitrophenyl(Dnp),
20 1-hydroxy-benzotriazole benzyl ester(OHbt), 8-hydroxy-quinoline ester(OHq), p-nitrobenzyloxy(ONBzl), phenylazophenyl and tertiary butoxy and is preferably Bzl.

X^6 is a protecting group for the side chain
25 guanidino group of Arg, such as nitro, Tos, trityl, benzyloxycarbonyl, adamantyloxycarbonyl, Z and Boc or X^6 may be hydrogen, which means there is no protection on the side chain group atoms. Tos is generally preferred.

30 X^7 may be Gly-NH-[resin support], D-Ala-NH-[resin support] or N(A)-[resin support]; or it may be amide either of Gly or of D-Ala or a substituted amide attached directly to Pro.

The criterion for selecting side chain
35 protecting groups for X^2 - X^6 is that the protecting group should be stable to the reagent under the reaction conditions selected for removing the α -amino protecting

group at each step of the synthesis. The protecting group should not be split off under coupling conditions, and the protecting group should be removable upon completion of the synthesis of the desired amino acid sequence under reaction conditions that will not alter the peptide chain.

When the X^7 group is Gly-NH-[resin support] or D-Ala-NH-[resin support], an amide bond connects Gly or D-Ala to BHA resin or to a MBHA resin. When the X^7 group is N(A)-[resin support], a substituted amide bond connects Pro to an N-alkylamino methyl resin (NAAM).

When X is acetyl, for example, at the N-terminus in the final formula, it may be possible to employ it as the X^1 protecting group for the α -amino group of D-NAL or whatever amino acid is used in the 1-position by adding it before the coupling of this last amino acid to the peptide chain. However, a reaction is preferably carried out with the peptide on the resin (after deblocking the α -amino group while the side-chain groups remain protected), e.g. by reacting with acetic acid in the presence of dicyclohexyl carbodiimide (DCC) or preferably with acetic anhydride or by another suitable reaction as known in the art.

The fully protected peptide intermediate results from classical solution synthesis and is then deprotected as is well known in the art. Deprotection of the peptide, as well as cleavage of the peptide from a BHA, MBHA or NAAM resin, is effected by treatment with hydrofluoric acid (HF) or its equivalent at a temperature which promotes formation of the acylium ion at the side-chain carboxyl group, preferably between about 20°C and about 25°C for an appropriate time, e.g. about 2-3 hours. A sufficient excess of a desired aromatic compound selected from Class Z', such as anisole, which also functions as a scavenger, is added to the peptide prior to treatment with HF. Generally an amount is added at least equal to 20 times the molar

amount of the peptide. In the presence of the acylium ion, this added compound from Class Z' reacts to create the aliphatic-aromatic ketone side chain, the mechanism being illustrated in Solid-Phase Peptide Synthesis, G. Barany & R. Merrifield, p. 192-197. After the removal of HF under vacuum, the cleaved, deprotected peptide is conveniently treated with ether, decanted, taken-up in dilute acetic acid and lyophilized. At this point, the peptide can, if desired, be converted to its nontoxic salt, as by treatment, for example, with 1 N acetic acid.

Broadly, the invention provides a method of making a peptide hormone of not greater than about fifty residues having a glutamic acid, a homoglutamic acid or an aspartic acid residue at a nonterminus position in the main chain thereof, the side chain of which residue constitutes a mixed alkyl ketone terminating in an aromatic group, which method comprises forming a peptide intermediate wherein said main peptide chain contains a glutamic acid, a homoglutamic acid or an aspartic acid residue in the desired position, the side chain carboxyl group of which is protected with a protecting group selected from the class consisting of Bzl(benzyl ester), 2,6-dichlorobenzyl(DCB), dinitrophenyl(Dnp), 1-hydroxy-benzotriazole benzyl ester(OHbt), 8-hydroxy-quinoline ester(OHq), p-nitrobenzyloxy(ONBzl), phenylazophenyl and tertiary butoxy; treating said peptide intermediate with HF and an aromatic compound selected from Class Z' (as defined herein) under conditions so that said protecting group is removed and an acylium ion intermediate is formed which ion reacts with said aromatic compound to form a mixed alkyl ketone therewith, and removing said HF and recovering said desired peptide hormone which has increased binding affinity to the receptor in question as a result of the inclusion of said aromatic ketone side chain. When making a GnRH nonapeptide or decapeptide, the residue is located in the 6-position.

More specifically, the invention provides a method for making a GnRH antagonist having Formula I or a nontoxic salt thereof, which method comprises (a) forming an intermediate compound having the Formula II:

5 $X^1-R_1-(W)D-Phe-R_3(X^2)-R_4(X^3)-R_5(X^4 \text{ or } X^6)-R_6(X^5)-R_7-Arg(X^6)-Pro-X^7$ wherein X^1 is hydrogen or an α -amino protecting group; X^2 is hydrogen or a protecting group for the indole nitrogen; X^3 is hydrogen or a protecting group for the alcoholic hydroxyl group of Ser or for a side-chain amino group; 10 X^4 is hydrogen or a protecting group for the phenolic hydroxyl group of Tyr; X^5 is a protecting group for a side chain carboxyl group, X^6 is hydrogen or a protecting group for a side-chain amino group; and X^7 15 is selected from the group consisting of Gly-NH-(resin support), D-Ala-NH-(resin support), -N(A)-(resin support), Gly-NH₂, D-Ala-NH₂, and substituted amides, wherein A represents an alkyl group; (b) splitting off one or more of the groups X^1 to X^6 20 and/or cleaving from any resin support included in X^7 by treatment with HF or its equivalent in an amount equal to about 5 to 15 times the weight of the resin plus a desired compound selected from Class Z' as defined hereinbefore and, if desired, (c) converting a 25 resulting peptide into a nontoxic salt thereof. The molar amount of the compound from Class Z' that is used is preferably at least about 50 times the molar amount of the synthetic peptide which is present.

Similar methods can be used for making GnRH 30 agonists and other peptide hormones of interest which are not more than about 50 residues long and which will exhibit increased binding affinity to the receptor in question as a result of the inclusion of such an aliphatic-aromatic ketone side chain on a nonterminal 35 residue.

Purification of the peptide is effected by ion exchange chromatography on a CMC column, followed by

partition chromatography using the elution system:
n-butanol;0.1N acetic acid (1:1 volume ratio) on a
column packed with Sephadex G-25, or by using HPLC, as
known in the art and reported in Rivier, J. et al.,

5 J. Chromatography, 288 (1984) 303-328.

The GnRH antagonists of the invention are
effective at levels of less than 100 micrograms per
kilogram of body weight, when administered at about noon
on the day of proestrous, to prevent ovulation in female
10 rats. For prolonged suppression of ovulation, it may be
necessary to use dosage levels in the range of from
about 0.1 to about 2.5 milligrams per kilogram of body
weight. These antagonists are also effective to arrest
spermatogenesis when administered to male mammals on a
15 regular basis and can thus be used as contraceptives.
Since these compounds will reduce testosterone levels
(an undesired consequence in the normal, sexually active
male), it may be reasonable to administer replacement
dosages of testosterone along with the GnRH antagonist.
20 These antagonists can also be used to regulate the
production of gonadotropins and sex steroids for other
purposes as indicated hereinbefore.

EXAMPLE I

GnRH antagonists as indicated in TABLE I having the formula:

Ac-R₁-(4Cl)D-Phe-R₃-Ser-Tyr-D-Glu(V)-Leu-Arg-Pro-R₁₀

5 are prepared by the solid-phase procedure referred to above, wherein Z' is a compound which results in the desired aromatic moiety portion V of the keto side chain.

TABLE I

	R ₁	R ₃	Z'	R ₁₀	
10					
	1	β-D-2NAL	D-3PAL	C ₆ H ₅ OCH ₃	D-Ala-NH ₂ , (Arg ⁵)
	2	"	D-Trp	"	"
	3	dehydro Pro	β-D-2NAL	"	Gly-NH ₂ , (4F)D-Phe ²
	4	β-D-2NAL	(6NH ₂)D-Trp	C ₆ H ₅ OH	"
15	5	"	(5OCH ₃)D-Trp	C ₆ H ₄ (OH) ₂	"
	6	"	(5Br)D-Trp	C ₆ H ₃ (CH ₃) ₂ OCH ₃	"
	7	"	(5F)D-Trp	C ₆ H ₅ SCH ₃	"
	8	"	(5Cl)D-Trp	C ₅ NH ₄ SH	"
	9	Pro	(5CH ₃)D-Trp	indole	D-Ala-NH ₂
20	10	β-D-2NAL	(N ⁱⁿ For)D-Trp	2-methylindole	"
	11	"	D-3PAL	3-methylindole	"
	12	Pro	(5Cl)D-Trp	"	"
	13	dehydro Pro	(6NO ₂)D-Trp	"	NHCH ₂ CH ₃
	14	D-Trp	(5F)D-Trp	"	"
25	15	D-pGlu	D-2PAL	"	D-Ala-NH ₂
	16	D-Phe	(6NO ₂)D-Trp	"	NHCH ₂ CH ₂ CH ₃

For purposes of an example, a representative solid phase synthesis of Peptide No. 1 above, which is referred to as [Ac-β-D-2NAL¹, (4Cl)D-Phe², D-3PAL³, Arg⁵, D-Glu⁶(C₆H₄OCH₃), D-Ala¹⁰]-GnRH is set forth hereinafter. This peptide has the following formula: Ac-β-D-2NAL-(4Cl)D-Phe-D-3PAL-Ser-Arg-D-Glu(C₆H₄OCH₃)-Leu-Arg-Pro-D-Ala-NH₂. The other peptides are similarly synthesized and purified.

A BHA resin is used, and Boc-protected D-Ala is coupled to the resin over a 2-hour period in CH₂Cl₂

using a 3-fold excess of Boc derivative and DCC as an activating reagent. The D-alanine residue attaches to the BHA residue by an amide bond.

Following the coupling of each amino acid residue, washing, deblocking and coupling of the next amino acid residue is carried out in accordance with the following schedule using an automated machine and beginning with about 5 grams of resin:

	STEP	REAGENTS AND OPERATIONS	MIX TIMES MIN.
10	1	CH ₂ Cl ₂ wash-80 ml. (2 times)	3
	2	Methanol (MeOH) wash-30 ml. (2 times)	3
	3	CH ₂ Cl ₂ wash-80 ml. (3 times)	3
	4	50 percent TFA plus 5 percent 1,2-ethanedithiol in CH ₂ Cl ₂ -70 ml. (2 times)	10
15	5	Isopropyl alcohol + 1% ethanedithiol wash-80 ml. (2 times)	3
	6	TEA 12.5 percent in CH ₂ Cl ₂ -70 ml. (2 times)	5
	7	MeOH wash-40 ml. (2 times)	2
20	8	CH ₂ Cl ₂ wash-80 ml. (3 times)	3
	9	Boc-amino acid (10 mmoles) in 30 ml. of either DMF or CH ₂ Cl ₂ , depending upon the solubility of the particular protected amino acid, (1 time) plus DCC (10 mmoles) in CH ₂ Cl ₂	30-300
25	10	MeOH wash-40 ml. (2 times)	3
	11	TEA 12.5 percent in CH ₂ Cl ₂ -70 ml. (1 time)	3
	12	MeOH wash-30 ml. (2 times)	3
30	13	CH ₂ Cl ₂ wash-80 ml. (2 times)	3

After step 13, if the synthesis is manual, an aliquot may be taken for a ninhydrin test: if the test is negative, go back to step 1 for coupling of the next amino acid; if the test is positive or slightly positive, go back and repeat steps 9 through 13.

The above schedule is generally used for coupling of each of the amino acids of the peptide of

the invention after the first amino acid has been attached. N^αBoc protection is used for each of the remaining amino acids throughout the synthesis. N^αBoc-β-D-2NAL is prepared by a method known in the art, 5 e.g. as described in detail in U.S. Patent No. 4,234,571, issued November 18, 1980. The side chain of Arg is protected with Tos. OBzl is used as a side chain protecting group for the hydroxyl group of Ser, and Bzl is used to protect D-Glu. D-3PAL is left unprotected. 10 N^αBoc-β-D-2NAL is introduced as the final amino acid. Boc-Arg(Tos), which has low solubility in CH₂Cl₂, is coupled using a DMF:CH₂Cl₂ mixture.

After deblocking the α-amino group at the N-terminal, its acetylation is achieved using a large 15 excess of acetic anhydride in dichloromethane. The cleavage of the peptide from the resin and complete deprotection of the side chains is carried out at 24°C. with HF for about 2-1/2 hours. A scavenger as set forth in the TABLE is added prior to HF treatment to produce 20 the mixed alkyl ketone. After the removal of HF under vacuum, the resin is extracted with 50% acetic acid, and the washings are lyophilized to provide a crude peptide powder.

Purification of the peptide is then effected by 25 ion exchange chromatography on CMC (Whatman CM 32, using a gradient of 0.05 to 0.3M NH₄OAc in 50/50 methanol/water) followed by partition chromatography in a gel filtration column using the elution system: n-Butanol; 0.1N Acetic acid (1:1 - volume ratio).

30 The peptide is judged to be homogeneous using thin layer chromatography and several different solvent systems, as well as by using reversed-phase high pressure liquid chromatography and an aqueous triethylammonium phosphate solution plus acetonitrile. 35 Amino acid analysis of the resultant, purified peptide is consistent with the formula for the prepared structure, showing substantially integer-values for each

amino acid in the chain. The optical rotation is **0192492** measured on a photoelectric polarimeter as $[\alpha]_D^{22} = -28.77^\circ \pm 1 (c=1, 50\% \text{ acetic acid})$.

The remaining GnRH antagonists set forth in
5 TABLE 1 are synthesized using the method specified above and an appropriate resin.

Each of the peptides is assayed in vivo to determine its effectiveness to prevent ovulation in female rats. In this test, a specified number of mature
10 female Sprague-Dawley rats, i.e. seven, each having a body weight from 225 to 250 grams, is injected subcutaneously with a specified microgram dosage of peptide in corn oil at about noon on the day of proestrous. Proestrous is the afternoon of ovulation.
15 A separate female rat group is used as a control to which the peptide is not administered. Each of the control female rats ovulates on the evening of proestrous; of the rats treated, the number of them which ovulate is recorded. Each of the peptides set
20 forth in Table I is considered to be significantly effective to prevent ovulation of female rats at a very low dosage, and each peptide is considered to be totally effective at a dose of about five micrograms.

EXAMPLE II

Peptides as indicated in TABLE II having the formula: Ac-B-D-2NAL-(W)D-Phe-D-Trp-R₄-R₅-R₆(V)-R₇-Arg-Pro-Gly-NH₂ are prepared by the solid-phase procedure referred to above, wherein Z' is employed to produce V.

TABLE II

	W	R ₄	R ₅	R ₆	Z'	R ₇
17	4F	Ser	Tyr	D-Glu	C ₆ H ₅ OCH ₃	Leu
10 18	4Br	"	(2F)Phe	D-Asp	"	Nle
19	"	AAL	Tyr	"	C ₆ H ₅ C ₇ H ₁₅	Nva
20	4Cl	aBu	"	D-Hgl	m-xylene	Nle
21	"	Ser	Arg	"	p-cymene	"
22	"	"	(2CH ₃)Phe	"	p-dibutylbenzene	Nva
15 23	4F	"	"	D-Asp	indane	NML
24	"	"	(3CH ₃)Phe	"	diethylindane	"
25	"	"	(2Cl)Phe	D-Glu	diphenylmethane	"
26	4NO ₂	"	Arg	"	C ₆ H ₅ (OH) ₂	"
27	"	Orn	Tyr	"	tetralin	Nle
20 28	2,4Cl ₂	Ser	(3F)Phe	"	iodobenzene	"
29	"	AAL	"	"	chlorophenol	Nva
30	C ^α Me/Cl	Ser	(3I)Tyr	"	1-phenyl-2-nitroethane	"
31	3,4Cl ₂	Orn	(3Cl)Phe	"	2-thiocresol	Leu

25

In vitro and/or in vivo testing of the peptides specified in Table II shows that the peptides listed in Table II are considered effective to block GnRH-induced LH secretion in vitro at a reasonable concentration.

30 Many of these peptides are more potent in vivo than the present standard. All of the peptides are considered to be effective to prevent ovulation of female mammals at very low dosages.

EXAMPLE III

Peptides as indicated in TABLE III having the formula: X-B-D-2NAL-(4Cl)D-Phe-(1 For)D-Trp-Ser-R₅-D-Glu(V)-NML-Arg-Pro-R₁₀ are prepared by the solid-phase procedure referred to above using an appropriate resin, wherein Z' is employed to produce V.

TABLE III

		X	R ₅	Z'	R ₁₀
10	32	Ac	Tyr	thioanisole	Gly-NH ₂
	33	Acr	"	diphenyl ether	D-Ala-NH ₂
	34	For	Arg	triethyl anisole	NHCH ₂ CH ₃
	35	Bz	(3F)Phe	chlorophenetole	NHCH ₃
	36	Ac	(2F)Phe	acetanilide	NHCF ₃
15	37	Vac	(2Cl)Phe	nitroanisole	NHCH ₂ CH ₂ CH ₃
	38	Acr	(3Cl)Phe	methyltolylether	NHCF ₂ CF ₃
	39	Ac	(3F)Phe	pyrogallol	D-Ala-NH ₂
	40	Acr	(3I)Tyr	salicylic acid	"
	41	Ac	Tyr	benzoic acid	"
20	42	"	(3Cl)Phe	benzoic acid	Gly-NH ₂
	43	Vac	"	biphenyl	NHNHCONH ₂
	44	Bz	Arg	diphenylbenzene	NHNHCONHCH ₃

In vitro and/or in vivo testing of the peptides specified in Table III shows that the peptides listed in Table III are considered effective to block GnRH-induced LH secretion in vitro at a reasonable concentration. Many of these peptides are more potent in vivo than the present standard. All of the peptides are considered to be effective to prevent ovulation of female mammals at very low dosages.

EXAMPLE IV

Peptides as indicated in TABLE IV having the formula: $\text{Ac-R}_1-(4\text{F})\text{D-Phe-R}_3\text{-Ser-Tyr-R}_6(\text{V})\text{-Leu-Arg-Pro-NHCH}_2\text{CH}_3$ are prepared by the solid-phase procedure referred to above, wherein Z' is used to produce V.

TABLE IV

		R_1	R_3	R_6	Z'
10	45	dehydro Pro	β -D-2NAL	D-Glu	methoxy biphenyl
	46	"	"	D-Asp	fluorene
	47	"	"	D-Hgl	anthracene
	48	"	"	D-Glu	phenanthrene
	49	β -D-1NAL	D-3PAL	D-Asp	indole-acetate salt
15	50	"	D-2PAL	D-Glu	furan
	51	Pro	"	"	methylbenzofuran
	52	D-Trp	"	D-Hgl	methyldibenzofuran
	53	D-Phe	"	D-Asp	chlorothiophene
	54	Pro	D-4PAL	D-Hgl	propyl pyrrole
20	55	"	"	"	acetyl carbazole
	56	D-pGlu	"	D-Glu	phenothiazine

In vitro and/or in vivo testing of the peptides specified in Table IV shows that the peptides listed in Table IV are considered effective to block GnRH-induced LH secretion in vitro at a reasonable concentration. Many of these peptides are more potent in vivo than the present standard. All of the peptides are considered to be effective to prevent ovulation of female mammals at very low dosages.

EXAMPLE V

GnRH agonists as indicated in TABLE V having the formula: pGlu-His-Trp-Ser-Tyr-R₆(V)-Leu-Arg-Pro-R₁₀ are prepared by the solid-phase procedure referred to 5 above, wherein Z' is used to produce V.

TABLE V

	R ₆	Z'	R ₁₀
57	D-Glu	C ₆ H ₄ OCH ₃	NHCH ₂ CH ₃
10 58	"	"	Gly-NH ₂
59	D-Asp	"	D-Ala-NH ₂
60	"	C ₆ H ₄ OH	Gly-NH ₂
61	D-Hgl	acridan	"
62	"	1,2-benzisoxazole	" (formyl Pro ¹)
15 63	"	phenothiazine	"
64	"	2-imidazolone	"
65	"	indole	D-Ala-NH ₂
66	"	1,2-dimethylindole	"
67	"	2-phenylindole	" (formyl Pro ¹)
20 68	D-Glu	tetrahydrocarbazole	"
69	"	hydroxyquinoline	NHCH ₂ CH ₃
70	D-Asp	resorcinol	"
71	"	phehnitene	D-Ala-NH ₂
72	D-Glu	durene	"
25			

To synthesize peptide No. 57, an N-ethylamine resin is used which is prepared by reacting a cross-linked chloromethylated polystyrene resin with ethylamine at 4°C. for about 2 days and Boc-protected 30 Pro is coupled to the resin over a 2-hour period in CH₂Cl₂ using a 3-fold excess of Boc derivative and DCC as an activating reagent. The proline residue attaches to the NEAM resin by a substituted amide bond.

Following the coupling of each amino acid 35 residue, washing, deblocking and coupling of the next amino acid residue is carried out in accordance with the following schedule using an automated machine and beginning with about 5 grams of resin:

STEP	REAGENTS AND OPERATIONS	MIX TIMES MIN.
1	CH ₂ Cl ₂ wash-80 ml. (2 times)	3
2	Methanol(MeOH) wash-30 ml. (2 times)	3
3	CH ₂ Cl ₂ wash-80 ml. (3 times)	3
5	4 50 percent TFA plus 5 percent 1,2-ethanedithiol in CH ₂ Cl ₂ -70 ml. (2 times)	10
5	5 Isopropyl alcohol + 1% ethanedithiol wash-80 ml. (2 times)	3
6	6 TEA 12.5 percent in CH ₂ Cl ₂ -70 ml. (2 times)	5
10	7 MeOH wash-40 ml. (2 times)	2
8	8 CH ₂ Cl ₂ wash-80 ml. (3 times)	3
9	9 Boc-amino acid (10 mmoles) in 30 ml. of either DMF or CH ₂ Cl ₂ , depending upon the solubility of the particular protected amino acid, (1 time)	30-300
15	10 MeOH wash-40 ml. (2 times)	3
11	11 TEA 12.5 percent in CH ₂ Cl ₂ -70 ml. (1 time)	3
20	12 MeOH wash-30 ml. (2 times)	3
13	13 CH ₂ Cl ₂ wash-80 ml. (2 times)	3

N^αBoc protection is used for each of the remaining amino acids throughout the synthesis, except for pGlu which is left unprotected; however, it can optionally be protected with Z. The side chain of Arg is protected with Tos. OBzl is used as a side chain protecting group for the hydroxyl group of Ser, and Bzl is used to protect D-Glu. Trp is left unprotected.

pGlu is introduced as the final amino acid. Boc-Arg(Tos) and Boc-Trp, which have low solubility in CH₂Cl₂, are coupled using DMF:CH₂Cl₂ mixtures.

After deblocking the α-amino group at the N-terminus, its acetylation is achieved using a large excess of acetic anhydride in dichloromethane. The cleavage of the peptide from the resin and complete deprotection of the side chains is carried out at 24°C.

with HF for about 2-1/2 hours. The compound Z' as set forth in the TABLE V is added prior to HF treatment to produce the mixed alkyl ketone and in most cases to act as a scavenger. After the removal of HF under vacuum, 5 the resin is extracted with 50% acetic acid, and the washings are lyophilized to provide a crude peptide powder.

Purification of the peptide is then effected by ion exchange chromatography on CMC (Whatman CM 32, using 10 a gradient of 0.05 to 0.3M NH_4OAc in 50/50 methanol/water) followed by partition chromatography in a gel filtration column using the elution system: n-Butanol; 0.1N Acetic acid (1:1 - volume ratio).

The peptide is judged to be homogeneous using 15 thin layer chromatography and several different solvent systems, as well as by using reversed-phase high pressure liquid chromatography and an aqueous triethylammonium phosphate solution plus acetonitrile. Amino acid analysis of the resultant, purified peptide 20 is consistent with the formula for the prepared structure, showing substantially integer-values for each amino acid in the chain.

The remaining GnRH agonists set forth in TABLE V are synthesized using the method specified above and an 25 appropriate resin.

Each of the peptides is assayed in vitro to determine its effectiveness to cause the secretion of LH from a primary culture of dispersed rat pituitary cells using the procedure set forth in U.S. patent No. 30 4,382,922. Each of the peptides set forth in Table V is considered to be very significantly more potent effective than native GnRH, and each peptide is considered to be totally effective at a reasonable dose to regulate fertility and to treat patients having 35 precocious puberty, endometriosis or dysmenorrhea.

The peptides of the invention are often administered in the form of pharmaceutically acceptable, nontoxic salts, such as acid addition salts, or of metal complexes, e.g., with zinc, barium, calcium, magnesium, 5 aluminum or the like (which are considered as addition salts for purposes of this application), or of combinations of the two. Illustrative of such acid addition salts are hydrochloride, hydrobromide, sulphate, phosphate, nitrate, oxalate, fumarate, 10 gluconate, tannate, maleate, acetate, citrate, benzoate, succinate, alginate, malate, ascorbate, tartrate and the like. An aqueous solution of the peptide is repeatedly treated, for example, with 1N acetic acid and then lyophilized to yield the acetic acid salt thereof. If 15 the active ingredient is to be administered in tablet form, the tablet may contain a pharmaceutically-acceptable diluent which includes a binder, such as tragacanth, corn starch or gelatin; a disintegrating agent, such as alginic acid; and a lubricant, such as 20 magnesium stearate. If administration in liquid form is desired, sweetening and/or flavoring may be used as part of the pharmaceutically-acceptable diluent, and intravenous administration in isotonic saline, phosphate buffer solutions or the like may be effected.

25 The pharmaceutical compositions will usually contain the peptide in conjunction with a conventional, pharmaceutically-acceptable carrier. Usually, the dosage will be from about 1 to about 100 micrograms of the peptide per kilogram of the body weight of the host 30 when given intravenously; oral dosages will be higher. Overall, treatment of subjects with these peptides is generally carried out in the same manner as the clinical treatment using other antagonists of GnRH.

These peptides can be administered to mammals 35 intravenously, subcutaneously, intramuscularly, orally, percutaneously, e.g. intranasally or intravaginally to achieve fertility inhibition and/or control and also in

applications calling for reversible suppression of gonadal activity, such as for the management of precocious puberty or during radiation or chemotherapy. Effective dosages will vary with the form of

5 administration and the particular species of mammal being treated. An example of one typical dosage form is a bacteriostatic water solution containing the peptide which solution is administered to provide a dose in the range of about 0.1 to 2.5 mg/kg of body weight. Oral
10 administration of the peptide may be given in either solid form or liquid form.

Although the invention has been described with regard to its preferred embodiments, it should be understood that changes and modifications as would be
15 obvious to one having the ordinary skill in this art may be made without departing from the scope of the invention which is set forth in the claims which are appended hereto. For example, other substitutions known in the art which do not significantly detract from the
20 effectiveness of the peptides may be employed in the peptides of the invention. For instance, instead of the residues specified for R_{10} , Sar-NH₂ (Sar = sarcosine) can be used, or NH-Y can be present, with Y being lower alkyl, cycloalkyl, fluoro lower alkyl or NHCONHQ, where
25 Q is H or lower alkyl, all of the foregoing being considered to be equivalents. D-Phe in the 1-position can be optionally halogenated as specified with respect to the 2-position substitution. In addition to the compounds enumerated as comprising Class 2', additional
30 equivalent aromatic compounds are identified in Volume 3, Aromatic Ketone Synthesis, Peter H. Gore, 1963, Interscience Publishers. Other equivalent residues, such as Met, Cys, Phe, Tyr and Trp can be used instead of those hereinbefore specified in the 7-position.
35 Particular features of the invention are emphasized in the claims that follow.

1. A peptide or a nontoxic salt thereof, said peptide having the formula: $X-R_1-R_2-R_3-R_4-R_5-R_6(V)-R_7-Arg-Pro-R_{10}$ wherein X is hydrogen or an acyl group having 7 or less carbon atoms; R_1 is pGlu, dehydro-Pro, Pro, D-pGlu, D-Phe, D-Trp or β -D-NAL; R_2 is (W)D-Phe or His; W is F, Cl, Cl_2 , Br, NO_2 or $C^{\alpha}Me/Cl$; R_3 is β -D-NAL, Trp, D-Trp, D-PAL, (N^{inFor})D-Trp or D-Trp which is substituted in the 5- or 6-position with NO_2 , NH_2 , OCH_3 , F, Cl, Br or CH_3 ; R_4 is Ser, Orn, AAL or aBu; R_5 is Tyr, Arg, (3F)Phe, (2F)Phe, (3I)Tyr, (3CH₃)Phe, (2CH₃)Phe, (3Cl)Phe or (2Cl)Phe; R_6 is D-Glu, D-Hgl or D-Asp; R_7 is Leu, NML, Nle or Nva; R_{10} is Gly-NH₂, D-Ala-NH₂ or NH-Y, with Y being lower alkyl, cycloalkyl, fluoro lower alkyl or NHCONHQ, where Q is H or lower alkyl; and V is an aromatic moiety portion of a ketone formed from the carboxylic group side chain of R_6 and a compound selected from Class Z' as defined herein.
2. A GnRH antagonist peptide or a nontoxic salt thereof according to Claim 1, wherein R_1 is dehydro-Pro, Pro, D-pGlu, D-Phe, D-Trp or β -D-NAL; and R_3 is D-Trp, D-3PAL, (N^{inFor})D-Trp or D-Trp which is substituted in the 5- or 6-position with NO_2 , NH_2 , OCH_3 , F, Cl, Br or CH_3 .
3. A peptide in accordance with Claim 2 wherein R_6 is D-Glu.
4. A peptide in accordance with Claim 2 or 3 wherein V is $C_6H_4OCH_3$.
5. A peptide in accordance with Claim 2 wherein R_6 is D-Hgl and V is $C_6H_4OCH_3$.
6. A peptide in accordance with Claim 2 having the formula: $Ac-\beta-D-2NAL-(4Cl)D-Phe-D-3PAL-Ser-Arg-D-Glu(C_6H_4OCH_3)-Leu-Arg-Pro-D-Ala-NH_2$.

7. A peptide in accordance with Claim 2 having the formula: Ac-dehydro Pro-(4F)D-Phe-β-D-2NAL-Ser-Tyr-D-Glu(C₆H₄OCH₃)-Leu-Arg-Pro-Gly-NH₂.

8. A GnRH agonist peptide or a nontoxic salt thereof according to Claim 1, said peptide having the formula: R₁-His-Trp-Ser-Tyr-R₆(V)-Leu-Arg-Pro-R₁₀ wherein R₁ is pGlu or formyl Pro and R₆ and R₁₀ are as defined therein.

9. A peptide in accordance with Claim 8 having the formula: pGlu-His-Trp-Ser-Tyr-D-Glu(C₆H₄OCH₃)-Leu-Arg-Pro-Gly-NHCH₂CH₃.

10. A method of making a GnRH analog peptide having a glutamic acid, a homoglutamic acid or an aspartic acid residue in the 6-position of the main chain thereof, the side chain of which residue constitutes a mixed alkyl ketone terminating in an aromatic group, which method comprises forming a nonapeptide or decapeptide intermediate wherein said main peptide chain contains a glutamic acid, a homoglutamic acid or an aspartic acid residue in the 6-position, the side chain carboxyl group of which is protected with a protecting group selected from the class consisting of benzyl ester, 2,6-dichlorobenzyl, dinitrophenyl, 1-hydroxy-benzotriazole benzyl ester, 8-hydroxy-quinoline ester, p-nitrobenzyloxy, phenylazophenyl and tertiary butoxy; treating said peptide intermediate with HF and an aromatic compound selected from Class Z' (as defined herein) under conditions so that said protecting group is removed and an acylium ion intermediate is formed which ion reacts with said aromatic compound to form a mixed alkyl ketone therewith, and removing said HF and recovering said desired GnRH analog peptide.

Claims for the contracting state: AT

1. A method of making a peptide hormone of not greater than about fifty residues having a glutamic acid, a homoglutamic acid or an aspartic acid residue at a non-terminus position in the main chain thereof, the side chain of which residue constitutes a mixed alkyl ketone terminating in an aromatic group, which method comprises forming a peptide intermediate wherein said main peptide chain contains a glutamic acid, a homoglutamic acid or an aspartic acid residue in the desired position, the side chain carboxyl group of which is protected with a protecting group selected from the class consisting of benzyl ester, 2,6-dichlorobenzyl, dinitrophenyl, 1-hydroxy-benzotriazole benzyl ester, 8-hydroxy-quinoline ester, p-nitrobenzyloxy, phenylazophenyl and tertiary butoxy; treating said peptide intermediate with HF and an aromatic compound selected from Class Z' (as defined herein) under conditions so that said protecting group is removed and an acylium ion intermediate is formed which ion reacts with said aromatic compound to form a mixed alkyl ketone therewith, and removing said HF and recovering said desired peptide hormone which has increased binding affinity to the receptor in question as a result of the inclusion of said aromatic ketone side chain.
2. A method according to Claim 1 for making a GnRH analog peptide having the formula: $X-R_1-R_2-R_3-R_4-R_5-R_6$ (M)- R_7 -Arg-Pro- R_{10} wherein X is hydrogen or an acyl group having 7 or less carbon atoms; R_1 is pGlu, dehydro-Pro, Pro, D-pGlu, D-Phe, D-Trp or β -D-NAL; R_2 is (W)D-Phe or His, W is F, Cl, Cl_2 , Br, NO_2 or $C^{\alpha}Me/Cl$; R_3 is β -D-NAL, D-Trp, Trp, D-PAL, (Nⁱⁿ For)D-Trp or D-Trp which is substituted in the 5- or 6-position with NO_2 , NH_2 , OCH_3 , F, Cl, Br or CH_3 ; R_4 is Ser, Orn, AAL or aBu; R_5 is Tyr, Arg, (3F)Phe, (2F)Phe, (3I)Tyr,

- (3CH₃)Phe, (2CH₃)Phe, (3Cl)Phe or (2Cl)Phe; R₆ is D-Glu, D-Hgl or D-Asp; R₇ is Leu, NML, Nle or Nva; R₁₀ is Gly-NH₂, D-Ala-NH₂ or NH-Y, with Y being lower alkyl, cycloalkyl, fluoro lower alkyl or NHCONHQ,
- 5 where Q is H or lower alkyl; and V is an aromatic moiety portion of a ketone formed from the carboxylic group side chain of R₆ and a compound selected from Class Z' as defined herein.
3. A method in accordance with Claim 2 wherein
- 10 R₆ is D-Glu.
4. A method in accordance with Claim 2 or 3 wherein V is C₆H₄OCH₃.
5. A method in accordance with Claim 2 wherein R₆ is D-Hgl and V is C₆H₄OCH₃.
- 15 6. A method in accordance with Claim 2 having the formula: Ac-β-D-2NAL-(4Cl)D-Phe-D-3PAL Ser-Arg-D-Glu(C₆H₄OCH₃)-Leu-Arg-Pro-D-Ala-NH₂.
7. A method in accordance with Claim 2 having the formula: Ac-dehydro Pro-(4F)D-Phe-β-D-2NAL-
- 20 Ser-Tyr-D-Glu(C₆H₄OCH₃)-Leu-Arg-Pro-Gly-NH₂.
8. A method according to Claim 1 for making a GnRH agonist peptide having the formula: R₁-His-Trp-Ser-Tyr-R₆(V)-Leu-Arg-Pro-R₁₀ wherein R₁ is pGlu or formyl Pro, R₆ is D-Glu, D-Hgl or
- 25 D-Asp; R₁₀ is Gly-NH₂, D-Ala-NH₂ or NH-Y, with Y being lower alkyl or fluoro lower alkyl; and V is an aromatic moiety portion of a ketone formed from a compound selected from Class Z' as defined herein.
9. A method in accordance with Claim 1 wherein
- 30 said peptide has the formula: pGlu-His-Trp-Ser-Tyr-D-Glu(C₆H₄OCH₃)-Leu-Arg-Pro-Gly-NHCH₂CH₃.
10. A method in accordance with Claim 1 wherein said treatment is carried out at a temperature of about 20°C. or above.

11. A method of contraceptive treatment of female mammals which method comprises administering to a female mammalian an amount of a GnRH antagonist effective to suppress or delay ovulation, characterized in 5 that said GnRH antagonist is a peptide as defined in Claim 2 (or a non-toxic salt thereof) in which R_2 is (W)D-Phe.

12. A method of contraceptive treatment of male mammals which method comprises administering to a male 10 mammalian an amount of a GnRH antagonist effective to suppress spermatogenesis, characterized in that said GnRH antagonist is a peptide as defined in Claim 2 (or a non-toxic salt thereof) in which R_2 is (W)D-Phe.

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☒ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)